

Reprinted from Chemosphere, Volume 32(3), Winters, D.; Cleverly, D.; Meier, K.; Dupuy, A.; Byrne, C.; Deyrup, C.; Ellis, R.; Ferrario, J.; Harless, R.; Leese, W.; Lorber, M.; McDaniel, D.; Schaum, J.; Walcott, J. 369-478. with permission from Elsevier Science.

The full article can be found in that reference. The abstract from this article is presented here.

A STATISTICAL SURVEY OF DIOXIN-LIKE COMPOUNDS IN UNITED STATES BEEF: A PROGRESS REPORT

Dwain Winters¹, David Cleverly², Kristen Meier⁴, Aubry Dupuy³, Christian Byrne³, Cindy Deyrup⁴, Richard Ellis⁴, Joseph Ferrario³, Robert Harless², William Leese⁴, Matthew Lorber², Danny McDaniel³, John Schaum², Jim Walcott⁴

¹Office of Prevention, Pesticides, and Toxic Substances, United States Environmental Protection Agency, Washington, D.C., 20460.

²Office of Research and Development, United States Environmental Protection Agency, Washington, D.C., 20460.

³Environmental Chemistry Laboratory, Office of Prevention, Pesticides, and Toxic Substances, United States Environmental Protection Agency, Stennis Space Center, Mississippi, 39529.

⁴Food Safety and Inspection Service, United States Department of Agriculture, Washington, D.C., 20250.

Abstract

The USEPA and the USDA have completed the first statistically designed survey of the occurrence and concentration of CDDs and CDFs in the fat of beef animals raised for human consumption in the United States. Back fat was sampled from 63 carcasses at federally inspected slaughter establishments nationwide. The sample design called for sampling beef animal classes in proportion to national annual slaughter statistics. All samples were analyzed using a modification of EPA method 1613, using isotope dilution, High Resolution GC/MS to determine the rate of occurrence of 2,3,7,8-substituted CDDs/CDFS. The method detection limits ranged from 0.05 ng kg⁻¹ for TCDD to 3 ng kg⁻¹ for OCDD. The results of this survey showed a mean concentration (reported as I-TEQ, lipid adjusted) in U.S. beef animals of 0.35 ng kg⁻¹ and 0.89 ng kg⁻¹ when either non-detects are treated as 0 value or assigned a value of 1/2 the detection limit, respectively.